


JIM AL-KHALILI

# QUANTUM



A Guide  
for the  
Perplexed

This book is dedicated to my father, to  
whom I am indebted for, among many  
things, first telling me about some  
strange theory called quantum mechanics.

First published in the United Kingdom in 2003  
by Weidenfeld & Nicolson

Published in the United States of America by Sterling Publishing Co., Inc  
387 Park Avenue South, New York, NY 10016-8810

Text copyright © Jim Al-Khalili, 2003  
Design and layout copyright © Weidenfeld & Nicolson, 2003

Pages 21-22: *The Road Not Taken* from *The Poetry of Robert Frost*  
edited by Edward Connery Lathem, published by Jonathan Cape.  
Reprinted by permission of The Random House Group Ltd.

The moral right of Jim Al-Khalili to be identified as the  
author of this work has been asserted in accordance  
with the Copyright, Design and Patents act of 1988.

All rights reserved. No part of this publication may be  
reproduced in any material form (including photocopying  
or storing it in any medium by electronic means and  
whether or not transiently or incidentally to some other  
use of this publication) without the written permission  
of the copyright owner, except in accordance with the  
provisions of the Copyright, Designs and Patents Act  
of 1988 or under the terms of a licence issued by the  
Copyright Licensing Agency, 90 Tottenham Court Road,  
London W1P 9HE. Applications for the copyright owner's  
written permission to reproduce any part of this publication  
should be addressed to the publisher.

Illustration: David Angel  
Design: Grade Design Consultants

No cats were harmed in the making of this book

A CIP catalogue record for this book is available from  
the British Library  
ISBN 0297843052

Printed and bound in Italy

Weidenfeld & Nicolson  
Wellington House  
125 Strand  
London WC2R 0BB

# Buckyballs and the Dual-Slit Experiment

*Professor Markus Arndt & Professor Anton Zeilinger* Department of Physics, University of Vienna

We usually think of a physical body as a localized object while the notion of a wave is intimately linked to something extended and delocalized. Contrary to this common belief quantum physics claims that both, seemingly contradictory, notions can apply to one and the same object in one and the same experiment.

We have recently implemented such an experiment with large carbon molecules called buckyballs. These molecules, known as  $C_{60}$  and  $C_{70}$ , contain sixty or seventy carbon atoms each, arranged to form the smallest known replica of a soccer ball, with a diameter no bigger than one millionth of a millimetre. In spite of their small size these molecules are the most massive objects ever used to demonstrate the wave-like nature of matter to date.

The experiment is set up as follows. The molecule source is a simple oven, filled with the carbon powder. The molecules can escape from a hole, like water vapour escaping from a hot kettle. They then fly through two collimating slits towards a laser detector with high resolution that can be shifted to record the spatial distribution of the molecular beam.

On the way towards the detector the molecules may encounter three different possibilities – either


no obstacle at all, or a very narrow slit or a very fine grating, which is a membrane with several slits.

The molecular beam profile for the first, 'empty', case is a single narrow peak and is in complete agreement with our naïve expectation, assuming that each molecule can be regarded as a free-flying classical ball.

However, the first weirdness occurs in the second case: If we place a single, very narrow slit – 70 nanometres (millionths of a millimetre) wide – in between the source and the detector we find a profile on the screen that differs from the empty case. We notice a strong broadening – instead of the narrowing, which we would have expected if the molecules were just little soccer balls. This is a consequence of diffraction, a property of waves.

The situation becomes even stranger when we replace the narrow slit by a grating. This structure is now composed of several openings, slightly narrower (nominally 50 nanometres) than the first slit. The slits are regularly spaced (about 50 nanometres apart). If molecules were simple particles we would expect an increased signal everywhere on the screen. But – to the surprise of our common sense – we now find that there are positions where we hardly detect any molecules at all.





Opening two or more pathways in the wall, instead of only one, reduces the number of detected molecules at certain places. This is very counter-intuitive and can no longer be explained with the model of classical balls flying along well defined paths, but it is in perfect agreement with a model based on the wave-nature of single molecules. Here we give up on the concept of a 'trajectory' and allow the molecules to simultaneously explore an extended space, which is orders of magnitude larger than the molecule itself, resulting in quantum interference.

It is important to note that the clicks in the detector are well localized and that the location where an individual molecule arrives is absolutely

random, as far as we can tell. But still, the weird wave-like pattern is built up as more and more molecules hit the detector.